

50X1-HUM

MAR 1952 51-4C

CLASSIFICATION CONFIDENTIALCENTRAL INTELLIGENCE AGENCY
INFORMATION FROM
FOREIGN DOCUMENTS OR RADIO BROADCASTS

REPORT

CD NO.

COUNTRY USSR
SUBJECT Scientific - Chemistry, toxic compoundsDATE OF
INFORMATION 1953HOW
PUBLISHED Monthly periodical

DATE DIST. 29 Jan 1954

WHERE
PUBLISHED Leningrad

NO. OF PAGES 2

DATE
PUBLISHED Aug 1953

LANGUAGE Russian

SUPPLEMENT TO
REPORT NO.

THIS DOCUMENT CONTAINS INFORMATION AFFECTING THE NATIONAL DEFENSE
OF THE UNITED STATES, WITHIN THE MEANING OF TITLE 18, SECTIONS 793
AND 794, OF THE U.S. CODE, AS AMENDED. ITS TRANSMISSION OR REVELA-
TION OF ITS CONTENTS TO OR RECEIPT BY AN UNAUTHORIZED PERSON IS
PROHIBITED BY LAW. THE REPRODUCTION OF THIS FORM IS PROHIBITED.

THIS IS UNEVALUATED INFORMATION


SOURCE Zhurnal Obshchey Khimii, Vol 23, No 8, pp 1431, 1432 α -GLYCOL ESTERS OF p-NITROPHENYLARSENOUS AND α -NAPHTHYLARSENOUS ACIDS

Gil'm Kamay and N. A. Chadayeva
Chem Inst im Acad A. Ye. Arbuzov
Kazan' Affiliate, Acad Sci USSR

[The experimental part of this article has been omitted.]

The present short work is a further development of our investigations in the field of cyclic glycol esters of arsenous and arylarsenous acids. Here we undertook the study of the methods of obtaining several cyclic α -glycol esters of p-nitrophenylarsenous and α -naphthylarsenous acids.

By the action of an equimolecular quantity of p-nitrophenylarsenous acid on various α -glycols (with heating), the α glycol esters of p-nitrophenylarsenous acid were synthesized according to the following scheme:



where: R = H, CH_2OCH_3 , or $\text{CH}_2\text{OC}_2\text{H}_5$.

- 1 -

CLASSIFICATION CONFIDENTIAL


STATE	NAVY	NSRB	DISTRIBUTION																
ARMY	AIR	FBI																	

50X1-HUM

CONFIDENTIAL

In this way, the ethyleneglycol, α -methoxypropyleneglycol, and α -ethoxypropyleneglycol esters of p-nitrophenylarsenous acid were prepared and then isolated. The esters readily saponify in water.

We prepared the α -glycol esters of α -naphthylarsenous acid by the reaction between α -naphthyldichloroarsine and the corresponding α -glycol in the presence of anhydrous pyridine. The reaction was carried out in absolute ether according to the following scheme:



The α -glycol esters of α -naphthylarsenous acid which are thus obtained are thick, transparent liquids unstable in air. They hydrolyze to form α -naphthylarsine oxide.

Conclusions: Various α -glycol esters of p-nitrophenylarsenous and α -naphthylarsenous acids have been synthesized and their properties studied.

BIBLIOGRAPHY

1. Gil'm Kamay and Z. L. Khisamova, DAN SSSR, Vol 46, p 535 (1951)
2. Gil'm Kamay and N. A. Chadayeva, DAN SSSR, Vol 81, p 71 (1952); Iz Ak Nauk SSSR, OTN, 908 (1952)
3. Gil'm Kamay, Z. L. Khisamova, and N. A. Chadayeva, DAN SSSR, Vol 89, 1051 (1953)

- E N D -

50X1-HUM

- 2 -

CONFIDENTIAL